arXiv:hep-th/9509076v1 14 Sep 1995

STATIC SPHERICALLY SYMMETRIC SOLUTIONS TO EINSTEIN-MAXWELL-DILATON FIELD EQUATIONS IN D DIMENSIONS

Metin Gürses and Emre Sermutlu
Department of Mathematics, Faculty of Science
Bilkent University, 06533 Ankara - Turkey
email:gurses@fen.bilkent.edu.tr

February 1, 2008

To appear in Classical and Quantum Gravity , 1995

Abstract

We classify the spherically symmetric solutions of the Einstein Maxwell Dilation field equations in D-dimensions and find some exact solutions of the string theory at all orders of the string tension parameter. We also show the uniqueness of the black hole solutions of this theory in static axially symmetric spacetimes.

1 Introduction

Recently we observe an increase of interest in constructing exact solutions of string theory [1]-[24]. There are several ways of such a construction [1]. One of these is to start with an exact solution of the leading order field equations (low energy limit of string theory) and showing that this solution solves also the string equations at all orders of the inverse power of the string tension parameter. Plane wave spacetimes with appropriate gauge and scalar fields have been shown to be exact solutions of the string theory [2]-[4]. In these solutions the higher order terms in the field equations dissapear due to the special form of the plane wave metrics. Lewi-Civita-Bertotti-Robinson spacetimes in four dimensions is also an exact solution of the string theory [8],[21]. In this solution the higher order terms do not dissapear but give some algebraic constraints among the constants of solutions [21].

To construct exact solutions in this approach one must first study the low energy limit of the string theory with some symmtery assumptions like the spherical symmetry. Among these solutions one searches for those satisfying the field equations at all orders. Hence we need a classification of the solutions under a symmetry of the low energy limit of the string theory. In this work ,instead of working directly with the low energy limit of the string theory we consider the Einstein Maxwell Dilaton field theory with arbitrary dilaton coupling constant α. For some fixed values of α this theory reduces to the low energy limit of the string theory and Kaluza Klein type of theories.

With the assumption of a symmetry , static spherical symmetry for instance , some of the solutions may describe black hole space times. Nonrotating black hole solutions of this theory , in spherically symmetric spacetimes , found so far [17]-[20] carry mass M , electric charge (or magnetic charge) Q and a scalar charge Σ. Out of these parameters only two of them are independent . In 4-dimensions we have the Garfinkle-Horowitz-Strominger (GHS) solution [18]. Gibbons and Maeda (GM) [17] have given black hole solutions in arbitrary D-dimensions. In four dimensions GM solution describes the region $r > r_1$ of the GHS metric. In this work we find and classify the solutions of the dilatonic Einstein-Maxwell (and hence of the low energy limit of the string theory) for static and spherically symmetric space-times.

Recently [22] the most general asymptotically flat spherically symmetric static solution of the low energy limit of string theory in four dimensions has been found. The uniqueness of the GHS-GM black hole solutions in four dimensional static spherically spacetimes is manifest in this work. Following this work we find the most general D dimensional spherically symmetric static solution of the Einstein Maxwell Dilaton field theory.

The uniquenes of these black hole solutions with the assumed symmetry (D dimensional spherical symmetry) is understood , becouse the solutions of the field equations are completely classified. On the other hand without solving the filed equations it is desirable to know the existance of other black hole solutions under relaxed symmetries. Here we show that GHS-GM metric in four dimensions is the unique asymptotically flat static axially symmetric black hole solution of the Einstein Maxwell Dilaton filed theory.

In this work we assume D dimensional static spherically symmetric spacetimes. We first write , in section 2 ,the geometrical quantitites like riemann and ricci tensors in terms of the volume form of the sphere S^{D-2} or in terms of its dual form. With such a representation it is relatively easier to work with the higher order curvature terms. In the third section we classify the static spherically symmetric solutions of the Einstein Maxwell Dilaton field equations. In the fourth section we prove that the GHS+GM metrics are the unique static axially symmetric black holes of the Einstein Maxwell Dilaton field theory. In the last section we give some theorems for the D dimesional static spherically symmetric spacetimes which are crucial to find the exact solutions of the string theory . In this section we give some possible exact solutions of string theory in D-dimensions.

2 Static Spherically Symmetric Riemannian Geometry

The line element of a static and spherically symmetric spacetime is given by:

$$ds^2 = -A^2 dt^2 + B^2 dr^2 + C^2 d\Omega^2_{D-2} \tag{1}$$

where A,B,C depend only on r. $d\Omega^2_{D-2}$ is the line element on S_{D-2}. The metric can be rewritten as $g_{ij} = -A^2 t_i t_j + B^2 k_i k_j + C^2 h_{ij}$, where $t_i = \delta^t_i$, $k_i = \delta^r_i$, h_{ij} =metric on D-2 sphere for $i, j \geq 2,\ h_{0i} = h_{1i} = 0$

Components of the christoffel symbol are given by

$$\begin{aligned} \Gamma^i_{jm} &= -AA'\, t^i (t_j k_m + t_m k_j) + AA'\, k^i t_j t_m + BB'\, k^i k_j k_m - CC'\, k^i h_{jm} \\ &\quad + CC'\, (h^i_j k_m + h^i_m k_j) + \Gamma^i_{(s)jm} \end{aligned} \tag{2}$$

where $\Gamma_{(s)}$ is the christoffel symbol on $D-2$ sphere

The Riemann tensor is given by

$$R^i_{jml} = \Gamma^i_{jl,m} - \Gamma^i_{jm,l} + \Gamma^i_{nm}\Gamma^n_{jl} - \Gamma^i_{nl}\Gamma^n_{jm} \tag{3}$$

We find that

$$\begin{aligned} R_{ijml} &= \left(AA'' - \frac{AA'B'}{B}\right) t_{[i}k_{j]}t_{[m}k_{l]} + \left(CC'' - \frac{CC'B'}{B}\right) k_{[i}h_{j][m}k_{l]} \\ &\quad + \frac{AA'CC'}{B^2}\, t_{[m}h_{l][j}t_{i]} - \frac{C^2 {C'}^2}{B^2} h_{i[m}h_{l]j} + C^2 R_{(s)ijml} \end{aligned} \tag{4}$$

where $R_{(s)ijml} = h_{i[m}h_{l]j}$ are the components of the Riemann tensor on S_{D-2}. Riemann tensor in (4) may be rewritten as

$$R_{ijml} = g_{jl}S_{im} - g_{jm}S_{il} + g_{im}S_{jl} - g_{il}S_{mj} + \eta_2 H_{ijk\cdots n}H_{ml}{}^{k\cdots n} \tag{5}$$

where

$$S_{ij} = \eta_0 M_{ij} + \eta_1 k_i k_j + \frac{1}{2}\eta_3 g_{ij} \tag{6}$$

$$M_{ij} = H_{im\cdots n}H_j^{m\cdots n} - \frac{1}{2(D-2)} H^2 g_{ij} \tag{7}$$

which turns out to be

$$M_{ij} = \frac{(D-3)!}{C^{2(D-3)}}\left(h_{ij} - \frac{1}{2C^2} g_{ij}\right) \tag{8}$$

Here the tensor $H_{ij\ldots k}$ is the volume form of S_{D-2}, i.e.

$$H_{ij\cdots k} = -\sqrt{h}\,\epsilon_{ij\cdots k} \tag{9}$$

where h denotes $\det h_{ij}$

Here the scalars η_i are given by

$$\eta_0 = \frac{C^{2(D-2)}}{(D-3)!}\left(\frac{A''}{AB^2} - \frac{A'B'}{AB^3} - \frac{C''}{CB^2} + \frac{B'C'}{B^3C}\right) \tag{10}$$

$$\eta_1 = \frac{A'C'}{AC} - \frac{C''}{C} + \frac{B'C'}{BC} \tag{11}$$

$$\eta_2 = \frac{C^{2(D-3)}}{(D-4)!}\left(1 - \frac{C'^2}{B^2} + \frac{A'C'C}{AB^2} - \frac{A''C^2}{AB^2} + \frac{A'B'C^2}{AB^3} + \frac{CC''}{B^2} - \frac{B'C'C}{B^3}\right) \tag{12}$$

$$\eta_3 = -\frac{A'C'}{AB^2C} \tag{13}$$

The Ricci tensor, Ricci scalar and Einstein tensor can be computed from Riemann as follows:

$$\begin{aligned} R_{ij} &= \left[\frac{(D-3)!}{2C^{2(D-2)}}(\eta_0(D-4)+\eta_2) + \frac{1}{B^2}\eta_1 + \eta_3(D-1)\right] g_{ij} \\ &\quad +\eta_1(D-2)k_ik_j + [\eta_0(D-2)+\eta_2]M_{ij} \end{aligned} \tag{14}$$

$$\begin{aligned} R &= \frac{(D-3)!}{C^{2(D-2)}}[\eta_0(D-4)(D-1)+\eta_2(D-2)] + \frac{2\eta_1(D-1)}{B^2} \\ &\quad +\eta_3 D(D-1) \end{aligned} \tag{15}$$

$$\begin{aligned}G_{ij} &= -\left[\frac{(D-3)!}{2C^{2(D-2)}}(\eta_0(D-4)(D-2)+\eta_2(D-3))+\frac{1}{B^2}\eta_1(D-2)\right. \\ &\left.+\frac{1}{2}\eta_3(D-1)(D-2)\right] g_{ij}+\eta_1(D-2)k_ik_j \\ &+[\eta_0(D-2)+\eta_2]M_{ij} \qquad (16)\end{aligned}$$

The covariant derivatives of $H_{ij\ldots k}$ and k_i are given as

$$\nabla_l H_{ij\ldots m} = -\rho[(D-2)H_{ij\ldots m}k_l + k_iH_{lj\ldots m} + k_jH_{il\ldots m} + \ldots + k_mH_{ij\ldots l}] \quad (17)$$

$$\nabla_i k_j = \rho_1 g_{ij} + \rho_2 M_{ij} + \rho_3 k_i k_j \quad (18)$$

where

$$\begin{aligned}\rho &= \frac{C'}{C} & \rho_1 &= \frac{A'C+AC'}{2AB^2C} \\ \rho_2 &= \frac{C^{2(D-2)}}{(D-3)!}\frac{A'C-AC'}{AB^2C} & \rho_3 &= -\frac{(AB)'}{AB} \qquad (19)\end{aligned}$$

In (5) we have given the Riemann tensor of a static, spherically symmetric spacetime in terms of the volume form of S_{D-2} (D-2)-brane field. We may also write the Riemann tensor in the form

$$R_{ijml} = g_{jl}S_{im} - g_{jm}S_{il} + g_{im}S_{jl} - g_{il}S_{mj} + e_2F_{ij}F_{ml} \quad (20)$$

where

$$F_{ij} = \frac{AB}{C^2}(t_ik_j - t_jk_i) \quad (21)$$

Other tensors are defined similar to the previous case, i.e.

$$S_{ij} = e_0\, \widetilde{M_{ij}} + e_1\, k_i k_j + \frac{1}{2} e_3\, g_{ij} \tag{22}$$

$$\widetilde{M_{ij}} = F_{mj} F_i^m - \frac{1}{4} F^2\, g_{ij} \tag{23}$$

$$\widetilde{M_{ij}} = \frac{1}{C^2} (h_{ij} - \frac{1}{2\, C^2}\, g_{ij}) \tag{24}$$

The scalars are given as

$$\begin{aligned} e_0 &= C^2 - \frac{{C'}^2 C^2}{B^2} + \frac{C^3 C' A'}{A\, B^2} \\ e_1 &= \eta_1 \\ e_2 &= -C^2 \left(1 - \frac{{C'}^2}{B^2} + \frac{A'C'C}{AB^2} - \frac{A''C^2}{AB^2} + \frac{A'B'C^2}{AB^3} + \frac{CC''}{B^2} - \frac{B'C'C}{B^3} \right) \\ e_3 &= \eta_3 \end{aligned} \tag{25}$$

Notice that when $D = 4$, $\widetilde{M_{ij}} = M_{ij}$, and $e_2 = -\eta_2$.

The covariant derivatives of F_{ij}, k_i and t_i are given as

$$\nabla_l F_{ij} = -\frac{3C'}{C} F_{ij} k_l - \frac{AC'}{BC^3} (t_j g_{il} - t_i g_{jl}) \tag{26}$$

$$\nabla_i k_j = \rho_1 g_{ij} + \tilde{\rho}_2 \widetilde{M_{ij}} + \rho_3 k_i k_j \tag{27}$$

$$\nabla_i t_j = \frac{A'}{A} (t_i k_j + t_j k_i) \tag{28}$$

where $\tilde{\rho}_2 = \frac{(D-3)!}{C^{2(D-4)}}\, \rho_2$

3 Solutions of the Einstein Maxwell Dilaton Field Equations

The field equations of the Einstein Maxwell Dilaton theory can be obtained from the following lagrangian

$$L = \sqrt{-g}\left[\frac{R}{2\kappa^2} - \frac{4}{(D-2)\kappa^2}(\nabla\phi)^2 - \frac{1}{4}\, e^{-\alpha_e \phi} F^2\right] \tag{29}$$

The field equations are

$$G_{ij} \;=\; \frac{8}{(D-2)}\left[\partial_i\phi\partial_j\phi - \frac{1}{2}(\nabla\phi)^2 g_{ij}\right] - \kappa^2 e^{-\alpha_e\phi}\left[F_i^{m}F_{jm} - \frac{1}{4}F^2 g_{ij}\right] \tag{30}$$

$$\nabla_i(e^{-\alpha_e\phi}F^{ij}) = 0, \tag{31}$$

$$\partial_i(\sqrt{-g}\, g^{ij}\,\partial_j\phi) + \frac{(D-2)\kappa^2\alpha_e\sqrt{-g}}{32}e^{-\alpha_e\phi}F^2 = 0, \tag{32}$$

where F_{ij} is the Maxwell and ϕ is the dilaton field. Here $i, j = 1, 2, ..., D \geq 4$. In static spherically symmetric spacetimes, gravitational field equations first lead to

$$\frac{\eta_0(D-4)(D-2)!}{2C^{2(D-2)}} + \frac{\eta_2(D-3)(D-3)!}{2C^{2(D-2)}} + \frac{\eta_1(D-2)}{B^2} + \frac{\eta_3(D-1)(D-2)}{2} - \frac{4\phi'^2}{(D-2)B^2} = 0 \tag{33}$$

$$\eta_1(D-2) - \frac{8\phi'^2}{(D-2)} = 0 \tag{34}$$

$$\eta_0(D-2)! + \eta_2(D-3)! - \frac{\kappa^2 Q^2 e^{\alpha_e \phi}}{A_{D-2}^2} = 0 \tag{35}$$

Dilaton equation is

$$\frac{8}{D-2}\left[\frac{A}{B}\, C^{D-2}\, \phi'\right]' - \frac{\alpha_e\, A\, B\, \kappa^2\, Q^2\, e^{\alpha_e \phi}}{2\, C^{D-2}\, A_{D-2}^2} = 0 \tag{36}$$

From (33) and (34) we obtain

$$\left[\frac{\left(A\, C^d\right)'\, C}{B}\right]' = d^2\, A\, B\, C^{d-1} \tag{37}$$

where $d = D-3$. Using the freedom in choosing the r coordinate we can let

$$A\, B\, C^{d-1} = r^{d-1}, \tag{38}$$

Using (37) and (38) we obtain

$$A^2\, C^{2d} = r^{2d} - 2b_1\, r^d + b_2 \tag{39}$$

where b_1 and b_2 are integration constants.

A combination of the dilaton (31) and gravitational field equations (30) give

$$d\, T^2 - \frac{d-1}{r}\, T + T' + \frac{8}{(d+1)^2}\, \phi'^2 = 0, \tag{40}$$

where T is defined as

$$T = \frac{(r^d + c_1)\, r^{d-1}}{(r^{2d} - 2b_1 r^d + b_2)} - \frac{16\phi'}{(d+1)^2 \alpha_e}. \tag{41}$$

Here c_1 is an integration constant Defining now a new function $\psi(\rho)$ as follows

$$\alpha_e\, \phi' = \frac{k_1\, d\, r^{d-1}}{(r^{2d} - 2b_1 r^d + b_2)}\, \psi(\rho) \tag{42}$$

the equation (40) becomes

$$\frac{r^{2d} - 2b_1 r^d + b_2}{d\, r^{d-1}}\, \frac{d\psi}{d\rho}\, \frac{d\rho}{dr} = (\psi + \mu)^2 + \nu^2 \tag{43}$$

The constants are given by

$$\begin{aligned} a &= \frac{(d+1)^2\, \alpha_e^2}{32\, d} \\ \mu &= -(c_1 + b_1) \\ \nu^2 &= a\mu^2 - \Delta(a+1) \end{aligned} \tag{44}$$

Now, if we solve the auxiliary equation

$$\frac{r^{2d} - 2b_1 r^d + b_2}{d\, r^{d-1}} \frac{d\rho}{dr} = \rho \tag{45}$$

and insert ρ above in Eq.(43) , we obtain

$$\rho \frac{d\psi}{d\rho} = (\psi + \mu)^2 + \nu^2, \tag{46}$$

Note that ρ depends on the sign of $\Delta = b_1^2 - b_2$. Hence ϕ can be found from ψ by

$$\phi = \frac{k_1}{\alpha_e} \int \frac{\psi(\rho)}{\rho} d\rho + \phi_0 \tag{47}$$

Where $k_1 = \frac{2\,a}{a+1}$ and ϕ_0 is an arbitrary constant. The metric function C is connected to ϕ as

$$\frac{C'}{C} = \frac{(r^d + c_1) r^{d-1}}{(r^{2d} - 2b_1 r^d + b_2)} - \frac{16\phi'}{(d+1)^2 \alpha_e} \tag{48}$$

This gives us

$$\ln(\frac{C}{c_0})^d = \left(\int \frac{u + b_1 + c_1}{u^2 - \Delta} du \right) - \frac{\alpha_e}{2a} \phi \tag{49}$$

Where $u = r^d - b_1$ and c_0 is an arbitrary integration constant.

Metric functions A and B can be found from C through the equations (38) and (39)

We have three different cases according to the sign of Δ.

$$\begin{array}{lll} \text{Case 1} & \Delta > 0 & \ln \rho(r) = \dfrac{1}{r_1^d - r_2^d} \ln \left(\dfrac{r^d - r_1^d}{r^d - r_2^d} \right) \\ \text{Case 2} & \Delta = 0 & \ln \rho(r) = -\dfrac{1}{r^d - r_3^d} \\ \text{Case 3} & \Delta < 0 & \ln \rho(r) = \dfrac{1}{\sqrt{-\Delta}} \left[\arctan \left(\dfrac{r^d - b_1}{\sqrt{-\Delta}} \right) - \dfrac{\pi}{2} \right] \end{array} \tag{50}$$

Where $r_1^d = b_1 + \sqrt{\Delta},\ r_2^d = b_1 - \sqrt{\Delta},\ r_3^d = b_1.$

The integration constants c_0 and ϕ_0 are determined by taking the asymptotic behaviour of the functions $C(r)$ and $\phi(r)$ to be

$$\lim_{r \to \infty} \phi = \phi_0 = 0 \tag{51}$$

$$\lim_{r\to\infty}\frac{C(r)}{r} = c_0 = 1 \tag{52}$$

To determine the remaining integration constants, we restrict the asymptotic behaviour of the metric, scalar field and the tensor field F_{ij} as follows:

$$\begin{aligned} \lim_{r\to\infty} r^d\,(A^2-1) &= -\frac{2\kappa^2\,M}{A_{d+1}\,(d+1)} \\ \lim_{r\to\infty} r^{d+1}\phi' &= -\frac{\kappa\sqrt{d+1}}{2A_{d+1}}\,\Sigma \\ \lim_{r\to\infty} r^{d+1}\,F_{tr} &= \frac{Q}{A_{d+1}} \end{aligned} \tag{53}$$

where M , Σ , and Q are the mass , dilaton and electric charges respectively [17]. From the definition of these constants it is clear that we are interested in the asymptotically flat solutions of the Einstein Maxwell Dilaton field equations

CASE 1

$$\Delta > 0 \tag{54}$$

Which means there are two roots to the equation (39). According to the sign of ν^2 we have three distinct solutions

Type 1

$$\begin{aligned} \nu^2 &< 0 \qquad\qquad \lambda^2 = -\nu^2 \\ \psi &= \frac{\lambda-\mu+c_2(\lambda+\mu)\rho^{2\lambda}}{1-c_2\rho^{2\lambda}} \end{aligned}$$

The metric functions become

$$
\begin{aligned}
A^2 &= \frac{(r^d - r_1^d)(r^d - r_2^d)}{C^{2d}} \qquad B^2 = \frac{r^{2d-2}\, C^2}{(r^d - r_1^d)(r^d - r_2^d)} && (55) \\
C^d &= (r^d - r_2^d) \left(\frac{1 - c_2\, \rho^{2\lambda}}{1 - c_2} \right)^{k_2} \rho^{k_3} && (56) \\
& && (57)
\end{aligned}
$$

Dilaton field is given as

$$
e^{\alpha_e\, \phi} = \left[\frac{(1 - c_2)\, \rho^{\lambda - \mu}}{1 - c_2\, \rho^{2\lambda}} \right]^{k_1} \tag{58}
$$

The constants are given by

$$
k_2 = \frac{1}{(a+1)}, \qquad k_3 = \frac{(r_1^d - r_2^d)}{2} - \frac{(\nu + a\mu)}{(a+1)} \tag{59}
$$

Undetermined integration constants are r_1, r_2, c_1 and c_2. From the boundary condition (53) we find that

$$
\begin{aligned}
2\, M &= \left[\frac{1 + c_2}{1 - c_2}\, \lambda + a\, \mu \right] e_1 \\
\Sigma &= \left[\frac{1 + c_2}{1 - c_2}\, \lambda - \mu \right] e_2 \\
Q &= \lambda\, \frac{\sqrt{c_2}}{1 - c_2} e_3
\end{aligned} \tag{60}
$$

The constants e_i are given by

$$
e_1 = \frac{2\, A_{d+1}\, (d+1)}{(a+1)\kappa^2}
$$

$$
\begin{aligned}
e_2 &= \frac{A_{d+1}\,\alpha_e\,(d+1)^{\frac{3}{2}}}{8\,(a+1)\kappa} \\
e_3 &= \frac{2\,A_{d+1}}{\kappa}\sqrt{\frac{(d+1)\,d}{(a+1)}}
\end{aligned}
\tag{61}
$$

Note that, we have four integration constants (c_1, c_2, r_1^d and r_2^d) but there exists only three equations to determine them. Also note that c_1 does not appear in the solution directly, so we have a freedom in c_1. ($c_2 \neq 1$) In order to complete the solution,we need to determine the integration constants in terms of the physical parameters M, Σ and Q. Let us define some auxiliary variables to solve the set of algebraic equations (60)

$$
\begin{aligned}
T_1 &= \frac{1}{a+1}(\frac{a\,\Sigma}{e_2} + \frac{2\,M}{e_1}) \qquad T_2 = \frac{1}{a+1}\,(\frac{2\,M}{e_1} - \frac{\Sigma}{e_2}) \\
T_3 &= \sqrt{1 - \frac{4\,Q^2}{e_3^2\,T_1^2}}
\end{aligned}
\tag{62}
$$

Then the integration constants are

$$
\begin{aligned}
\lambda &= T_3 T_1 \\
\mu &= T_2 \\
c_2 &= \frac{1-T_3}{1+T_3}
\end{aligned}
\tag{63}
$$

Also

$$
\Delta = \frac{a\mu^2 + \lambda^2}{a+1} \tag{64}
$$

The reality of T_3 imposes

$$M + g\,\Sigma \geq s\,|Q| \tag{65}$$

where g and s are given by

$$g = \frac{\alpha_e\,(d+1)^{3/2}}{4d\,\kappa} \quad , \quad s = \frac{1}{\kappa}\sqrt{\frac{(d+1)(a+1)}{d}} \tag{66}$$

Such an inequality has been found by Gibbons and Wells for D=4.

When $\Delta > 0$ we have two roots to the equation (39). In general these roots are the singular points of the space-time. If the integration constants satisfy some additional constraints one of these roots becomes regular. An invariant of the space-time is the scalar curvature is given by

$$R = A_1 \frac{\rho^{z_1}\left[-\mu+\lambda+\frac{2c_2\lambda\rho^{2\lambda}}{1-c_2\rho^{2\lambda}}\right]^2}{[(r^d-r_1^d)(r^d-r_2^d)]^{1+\frac{1}{d}}(1-c_2\rho^{2\lambda})^{z_2}} + A_2 \frac{\rho^{z_1+2\lambda}\left(\frac{1-c_2}{1-c_2\rho^{2\lambda}}\right)^{z_2+2}}{[(r^d-r_1^d)(r^d-r_2^d)]^{1+\frac{1}{d}}} \tag{67}$$

where

$$\begin{aligned} z_1 &= \frac{2}{d(a+1)}(a\mu+\lambda) \\ z_2 &= \frac{2}{d(a+1)} \\ A_1 &= \frac{8}{d+1}\frac{k_1^2 d^2}{\alpha_e^2} \\ A_2 &= \frac{\kappa^2}{2}\frac{(D-4)}{(D-2)} \end{aligned} \tag{68}$$

As $r \longrightarrow r_1$ we have a singularity unless we choose $\mu = \lambda$. This choice, by the utility of eqn (44), gives $\mu = \lambda = \frac{r_1^d - r_2^d}{2}$. Inserting this in equation (67), we find that as $r \longrightarrow r_1$,

$$R = \tilde{A}_1\,(r^d - r_1^d) + \tilde{A}_2 \tag{69}$$

where $\tilde{A}_1$ and $\tilde{A}_2$ are constants , so the horizon is regular. The choice $\mu = \lambda$ gives $T_2 = T_1 T_3$ by equaton (63), which means

$$Q^2 = \frac{8d\Sigma}{\alpha_e}\left[\frac{2\kappa}{(d+1)^{\frac{3}{2}}}M + \frac{8(a-1)}{\alpha_e(d+1)^2}\Sigma\right] \tag{70}$$

The connection between physical parameters and integration constants is

$$\begin{aligned} 2M &= \frac{r_1^d - r_2^d}{2}\left(\frac{1+c_2}{1-c_2} + a\right)e_1 \\ \Sigma &= \frac{r_1^d - r_2^d}{2}\frac{2c_2}{1-c_2}e_2 \qquad (71) \\ Q &= \frac{r_1^d - r_2^d}{2}\frac{\sqrt{c_2}}{1-c_2}e_3 \end{aligned}$$

$$(72)$$

The metric is

$$ds^2 = -\frac{(r^d - r_1^d)(r^d - r_2^d)}{C^{2d}}dt^2 + \frac{r^{2d-2}C^2}{(r^d - r_1^d)(r^d - r_2^d)}dr^2 + C^2 d\Omega_{D-2}^2 \tag{73}$$

where

$$C^d = (r^d - r_2^d)\left(1 + \frac{\Sigma/e_2}{r^d - r_2^d}\right)^{k_2} \tag{74}$$

$$e^{\alpha_e\,\phi} = \left(1 + \frac{\Sigma/e_2}{r^d - r_2^d}\right)^{-k_1} \tag{75}$$

$$F_{tr} = \frac{Q}{A_{d+1} r^{d+1}} \tag{76}$$

and $\frac{r_1^d - r_2^d}{2} = T_2$.

If we choose the integration constant r_2 as zero, these black hole metrics are identical with the metrics given by Gibbons and Maeda [17]
Although the extreme limit ($\Delta = 0$) in genereal is going to be studied in Case II we obtain the extreme case of the above solution by setting $c_2 = (r_2/r_1)^d$ and $r_2 = r_1$. Then we obtain

$$\begin{aligned} 2M &= r_1^d e_1 \\ \Sigma &= r_1^d e_2 \qquad (77) \\ 2Q &= r_1^d e_3 \end{aligned}$$

$$\tag{78}$$

Type 2

$$\begin{aligned} \nu^2 &> 0 \\ \psi &= \nu \, \tan(c_2 + \nu \ln \rho) - \mu \end{aligned} \tag{79}$$

$$\int \frac{\psi(\rho)}{\rho} d\rho = -\ln\left[\cos(c_2 + \nu \ln \rho)\right] - \mu \ln \rho + c_3 \tag{80}$$

After similar steps as the previous type, we arrive at the solution

$$A^2 = \frac{(r^d - r_1^d)(r^d - r_2^d)}{C^{2d}} \qquad B^2 = \frac{r^{2d-2}\, C^2}{(r^d - r_1^d)(r^d - r_2^d)} \tag{81}$$

$$C^d = (r^d - r_2^d)\, \rho^{\frac{r_1^d - r_2^d}{2} - \frac{\mu\, k_1}{2}} \left[\frac{\cos(c_2 + \nu \ln \rho)}{\cos c_2}\right]^{k_2} \tag{82}$$

Scalar field is given as

$$e^{\alpha_e\, \phi} = \left[\frac{\cos c_2}{\rho^{\mu}\, \cos(c_2 + \nu \ln \rho)}\right]^{k_1} \tag{83}$$

Physical parameters are found using (53)

$$\begin{aligned} 2M &= (a\mu + \nu \tan c_2)\, e_1 \\ \Sigma &= (-\mu + \nu \tan c_2)\, e_2 \\ Q &= \frac{\nu}{\cos c_2}\, e_3 \end{aligned} \tag{84}$$

$$\begin{aligned} \mu &= T_2 \\ \sin c_2 &= \frac{e_3\, T_1}{Q} \\ \nu \tan c_2 &= T_1 \end{aligned} \tag{85}$$

The condition $|\sin c_2| < 1$ imposes

$$M + g\,\Sigma < s\,|Q| \tag{86}$$

Where g and s are defined in (66). We also have to check the sign of Δ.

$$(a+1)\Delta = \frac{1}{a+1}(\frac{a\,\Sigma^2}{e_2^2} + \frac{4\,M^2}{e_1^2}) - \frac{Q^2}{e_3^2} \tag{87}$$

Here the sign of Δ puts a constraint on the physical variables.

Type 3

$$\begin{aligned} \nu^2 &= 0 \\ \psi &= -\mu - \frac{1}{\ln\rho + c_2} \end{aligned} \tag{88}$$

$$\int \frac{\psi(\rho)}{\rho} d\rho = -\ln(\ln\rho + c_2) - \mu\ln\rho + c_3 \tag{89}$$

$$e^{\alpha_e\phi} = \left[\frac{c_2}{\rho^\mu(c_2+\ln\rho)}\right]^{k_1} \tag{90}$$

$$C^d = \left(\frac{c_2+\ln\rho}{c_2}\right)^{k_2} [(r^d - r_1^d)(r^d - r_2^d)]^{\frac{1}{2}} \rho^{\frac{-\mu k_1}{2}} \tag{91}$$

Physical parameters are found using (53)

$$\begin{aligned} 2M &= (a\mu - \frac{1}{c_2})\,e_1 \\ \Sigma &= (-\mu - \frac{1}{c_2})\,e_2 \\ Q &= \frac{e_3}{2c_2} \end{aligned} \tag{92}$$

The solution is

$$\begin{aligned} \mu &= -T_2 \\ -\frac{1}{c_2} &= T_1 \qquad (93) \\ & \qquad (94) \end{aligned}$$

which gives

$$\frac{2\,Q}{e_3} = T_1 \tag{95}$$

This is the equality case of the inequality (65)

CASE 2

$$\Delta = 0 \tag{96}$$

Then there is one root to the equation (39). Denote it by r_1^d.

$$A\,C^d = (r^d - r_1^d) \tag{97}$$

$$A^2 = \frac{(r^d - r_1^d)^2}{C^{2d}} \qquad B^2 = \frac{r^{2d-2}\,C^2}{(r^d - r_1^d)^2} \tag{98}$$

$$C^d = (r^d - r_1^d)\,\rho^{-\frac{\mu\,k_1}{2}} \left[\frac{\cos(c_2 + \nu \ln \rho)}{\cos(c_2)}\right]^{k_2} \tag{99}$$

Scalar field is given as

$$e^{\alpha_e\,\phi} = \left[\frac{\cos c_2}{\rho^{\mu}\ \cos(c_2 + \nu \ln \rho)}\right]^{k_1} \tag{100}$$

CASE 3

$$\Delta < 0 \tag{101}$$

This case is similar to the previous one.

$$e^{\alpha_e \phi} = \left[\frac{\cos c_2}{\rho^{\mu} \, \cos(c_2 + \nu \ln \rho)} \right]^{k_1} \tag{102}$$

$$A^2 = \frac{(r^{2d} - 2b_1 r^d + b_2)}{C^{2d}} \qquad B^2 = \frac{r^{2d-2} \, C^2}{(r^{2d} - 2b_1 r^d + b_2)} \tag{103}$$

$$C^d = (r^{2d} - 2b_1 r^d + b_2)^{\frac{1}{2}} \, \rho^{-\frac{\mu \, k_1}{2}} \left[\frac{\cos(c_2 + \nu \ln \rho)}{\cos(c_2)} \right]^{k_2} \tag{104}$$

In cases 2 and 3, the relation of physical parameters to integration constants are exactly the same as case 1 type 2. In addition we have the equation (87) with the sign of Δ chosen according to the case.

It is perhaps noticed that we solve field equations exactly without assuming any special ansatz [1] . In that respect we have all the possible integration constants in the solutions , hence we have the following assertion: In D-dimensions the metric given in (73) and (74) is the only asymptotically flat metric corresponding to a spherically symmetric static black hole carrying mass M , dilaton charge Σ and electric charge Q and covered by a regular horizon. In D-dimensions the Gibbons-Maeda metric is diffeomorphic to the region $r > r_1$ where r_1 is the location of the outer horizon. Relaxing the condition of asymptotical flatness there is a possibility having a black hole

---

[1] After this work was completed we became aware of a recent paper [23] , which similarly solves the field equations (30)-(32) without any special ansatz, but in completely different to those used here

(with a regular horizon and singularity located at the origin) solution of the Einstein Maxwell Dilaton field equations [24]

Expressing the boundary conditions of the black hole solution (74) (the behaviour of the metric as r approaches to the outer horizon r_1 and to infinity) as the black hole boundary conditions in static axially symmetric four dimensions one can prove that GHS or Gibbons-Maeda solution describes also the unique asymptotically flat static black hole solution in Einstein Maxwell Dilaton Gravity. In the next section we give the proof of this statement.

4 Uniqueness of Black Hole Soltions

In this section we are interested in the black hole solutions of the Einstein Maxwell Dilaton theory in the static axially symmteric spacetimes in four dimensions. Using a different approach the uniqueness of static charged dilaton black hole (GHS+GM black hole) has been recently shown in ref.[27]. In this proof the dilaton coupling constant was taken fixed ($a = 2$) which corresponds to the low energy limit of the string theory. Here we shall show that the static black holes of the Einstein Maxwell Dilaton theory are unique for arbitrary values of the dilaton coupling constant a. We are not going to solve the corresponding field equations but first formulate these field equations as a sigma model in two dimensions and use this formulation in the proof of the uniqueness of the solutions under the same boundary condtions. Uniqueness of the stationary black hole solutions of the Einstein theory is now a very well established concept [27]-[34]. This proof is based on the sigma model formulation of the stationary axially symmteric Einstein Maxwell field equations

[30]-[37]. Here we shall follow the approach given by [30] and [34].

The line element of a static axially symmetric four dimensional spacetime is given by

$$ds^2 = e^{2\,\psi}\,[e^{2\gamma}\,(d\rho^2 + dz^2) + \rho^2\,d\phi^2\,] - e^{-2\,\psi}\,dt^2, \tag{105}$$

The field equations of the Einstein Maxwell Dilaton field theory with the above metric and $A_\mu = (A, 0, 0, 0)$ are

$$\nabla^2\,\psi + \frac{\kappa^2}{2}\,e^{2\,\psi - a\,\phi}\,(\nabla\,A)^2 = 0 \tag{106}$$

$$\nabla^2\,\phi - a\,\frac{\kappa^2}{8}\,e^{2\,\psi - a\,\phi}\,(\nabla\,A)^2 = 0 \tag{107}$$

$$\nabla^2\,A + \nabla(2\psi - a\,\phi)\,\nabla\,A = 0 \tag{108}$$

$$\frac{1}{\rho}\,\gamma_{,z} = 2\,\psi_{,\rho}\,\psi_{,z} + 4\,\phi_{\rho}\,\phi_{,z} - \kappa^2\,e^{2\,\psi - a\,\phi}\,A_{\rho}\,A_{,z} \tag{109}$$

$$\frac{2}{\rho}\,\gamma_{,\rho} = 2\,\psi^2_{,\rho} - 2\,\psi^2_{,z} + 4\,\phi^2_{,\rho} - 4\,\phi^2_{,z} - \kappa^2\,e^{2\,\psi - a\,\phi}\,(A^2_{,\rho} -\,A^2_{,z}) \tag{110}$$

let $E = \psi - \frac{a}{2}\phi$ and $B = \frac{\kappa'}{\sqrt{2}}\,A$ we then find

$$\nabla^2\,E +\,e^{2\,E}\,(\nabla\,B)^2 = 0 \tag{111}$$

$$\nabla^2\,B + 2\,\nabla\,E\,\nabla\,B = 0 \tag{112}$$

where

$$\kappa' = \kappa \sqrt{1 + \frac{a^2}{8}} \tag{113}$$

We wish to write the Eqs.(111 , 112) as a single complex equation by introducing a complex potential. In order to achieve this we introduce pseudopotential ω by use of (112)

$$\omega_\rho = \rho\, e^{-2E}\, B_z \quad , \quad \omega_z = -\rho\, e^{-2E}\, B_\rho \tag{114}$$

then the resulting equations can be written as the following single complex equation (the Ernst equation) for $\varepsilon = \rho\, e^E + i\omega$

$$Re(\varepsilon)\, \nabla^2\, \varepsilon = \nabla\, \varepsilon\, \nabla\, \varepsilon \tag{115}$$

Hence the above complex equation reperesents the Eqs.(111) and (112) if we let $\varepsilon = \rho\, e^E + i\, \omega$. The above Ernst equation (115) defines a sigma model on $SU(2)/U(1)$ with the equation of motion

$$\vec{\nabla}\, \left(P^{-1}\, \vec{\nabla}\, P\right) = 0 \tag{116}$$

where

$$P = \frac{1}{\rho e^E} \begin{pmatrix} 1 & -B \\ -B & \rho^2 e^{2E} + B^2 \end{pmatrix} \tag{117}$$

In the sequel we assume enough differentiability for the components of the matrix P in $V \cup \partial V$. Here V is a region in M with boundary ∂V. In our case V is the region $r > 0$ (see section 2 type I solutions) and hence ∂V has two disconnected components.

We also assume that P is positive definite. Let P_1 and P_2 be two different solutions of (116). The difference of their equations satisfy

$$\vec{\nabla}\left(P_1^{-1}(\vec{\nabla}\, Q)\, P_2\right) = 0. \tag{118}$$

where $Q = P_1\, P_2^{-1}$. Multiplying the both sides by $Q^\dagger$ (hermitian conjugation) and taking the trace we obtain

$$\vec{\nabla}\left[tr\left(Q^\dagger\, P_1^{-1}(\vec{\nabla}Q)\, P_2\right)\right] = tr\left[(\vec{\nabla}Q^\dagger)\, P_1^{-1}\,(\vec{\nabla}Q)\, P_2\right] \tag{119}$$

The left hand side of the above equation can be simplified further and we obtain

$$\nabla^2\, q = tr\left[(\vec{\nabla}Q^\dagger)\, P_1^{-1}(\vec{\nabla}Q)\, P_2\right] \tag{120}$$

n where $q = tr(Q)$. Using the hermiticity and positive definiteness properties of the matrices P_1 and P_2 we may let

$$P_i = A_i\, A_i^\dagger,\ \ (i = 1, 2) \tag{121}$$

where A_1 and A_2 are nonsingular $n \times n$ matrices given by

$$A_i = \frac{1}{\sqrt{\rho}e^{\frac{E_i}{2}}}\begin{pmatrix} 1 & 0 \\ -B_i & \rho e^{E_i} \end{pmatrix} \tag{122}$$

With the aid of (121) Eq.(120) reduces to

$$\nabla^2 q = tr(\vec{J}^\dagger \, \vec{J}) \tag{123}$$

where

$$\vec{J} = A_1^{-1} (\vec{\nabla} Q) \, A_2 \tag{124}$$

Eq.(123) is a crucial step towards the proof of the uniqueness theorems. It is clear that the right hand side is positive definite at all point of V. Before going on let us give the scalar function q.

$$q = 2 + \frac{1}{\rho^2 \, e^{E_1 + E_2}} \left[\rho^2 \, (e^{E_1} - e^{E_2})^2 \, + \, (B_1 \, - \, B_2)^2 \right] \tag{125}$$

It is clear that $q = 2$ and its first derivatives vanish on the boundary ∂V of V.

Let M be an 2 dimensional manifold with local coordinates (ρ, z). Let V be a region in M with boundary ∂V. Let P be a hermitian positive definite 2×2 matrix with unit determinant and let P_1 and P_2 be two such matrices satisfying (116) in V with the same boundary conditions on ∂V then we have $P_1 = P_2$ at all points in region V . The proof is as follows. Integrating (123) in V we obtain

$$\int_{\partial V} \vec{\nabla} \, q \, d \, \vec{\sigma} = \int_V tr \, (\vec{J}^\dagger \, \vec{J}) \, dV \tag{126}$$

and using the boundary condition $q = 2$ on ∂V we get

$$\int_V tr \, (\vec{J}^\dagger \, \vec{J}) \, dV = 0 \tag{127}$$

Then the integrand in (127) vanishes at all points in V. This implies the vanishing of $\vec{J}$ which implies that $Q = Q_0 =$ a constant matrix in V. Since Q is the identity matrix I on ∂V then $Q = I$ in V. Hence $P_1 = P_2$ at all points in V. Another way to obtain this result is to use (123) directly. The vanishing of the integrand in (127) implies that q is an harmonic function in V. Since $q = 2$ on the boundary ∂V of V then it must be equal to the same constant in V as well. This implies that $P_1 = P_2$ in V.

In four dimensions the Einstein Maxwell Dilaton field theory a static black hole should carry mass M , electric charge Q and dilaton charge Σ. Such a black hole solution exist which was found by Gibbons and Maeda for an arbitrary dilaton coupling parameter a. Here the above proof implies that all those solutions with the same black hole boundary conditions (asymptotically flat and regular horizons) as the GM solution are the same everywhere in spacetime.

5 Exact Solutions of String Theory

In this section we introduce some theorems for static spherically symmetric D dimesional spacetimes which will be utilized to obtain exact solutions of string theory. This section is a natural extension of the work in ref.([21]) to an arbitrary dimensions. We shall not give the proofs of the theorems. The necessary tools for the proofs are given in the second section.

The covariant derivatives of $H_{ij\cdots k}$ and k_i given in (17) and (18) are expressed only in terms of themselves and the metric tensor. Hence we have the following theorem:

Theorem 1 *Covariant derivatives of the Riemann tensor* R_{ijkl}*, the tensor* $H_{ij\cdots k}$ *and the vector* k_i *at any order are expressible only in terms of* $H_{ij\cdots k}$ *,* g_{ij} *,* k_i

Since contraction of k^i with $H_{ij\cdots k}$ vanishes, the only symmetric tensors constructable out of $H_{ij\cdots k}$, g_{ij} and k_i are M_{ij}, the metric tensor g_{ij} and k_ik_j. Then the following theorem holds:

Theorem 2 *Any second rank symmetric tensor constructed out of the Riemann tensor, anti-symmetric tensor* $H_{ij\cdots k}$*, dilaton field* $\phi = \phi(r)$ *and their covariant derivatives is a linear combination of* M_{ij} *,* g_{ij} *and* k_ik_j*.*

Let this symmetric tensor be E'_{ij}. Then we have

$$E'_{ij} = \sigma_1 M_{ij} + \sigma_2 g_{ij} + \sigma_3 k_i k_j \tag{128}$$

where σ_1, σ_2 and σ_3 are scalars which are functions of the metric functions, invariants constructed out of the curvature tensor R_{ijkl}, $H_{ij\cdots k}$ and the dilaton field $\phi(r)$ and their covariant derivatives.

Theorem 3 *Any vector constructed out of the Riemannian tensor* R_{ijkl}*,* $H_{ij...k}$ *the dilaton field* $\phi = \phi(r)$ *and their covariant derivatives is proportional to* k_i*.*

Let this vector be E'_i. Hence

$$E'_i = \sigma k_i \tag{129}$$

where σ is a scalar like $\sigma_1,\ \sigma_2,\ \sigma_3$.

The covariant derivatives of F_{ij}, k_i and t_i are expressed in terms of themselves, metric tensor. So we have a similar theorem:

Theorem 4 *Covariant derivatives of the Riemann tensor* R_{ijkl}*, the anti-*

symmetric tensor F_{ij}, the vectors k_i and t_i at any order are expressible only in terms of F_{ij}, k_i, t_i and g_{ij}.

As an example to an exact solution of the low energy limit of the string theory in D-dimensions with constant dilaton field, we propose the LCBR (Levi-Civita Bertotti-Robinson) metric which is given by

$$g_{ij} = \frac{q^2}{r^2}\,(-t_i t_j + c_0^2 k_i k_j - r^2 h_{ij}) \tag{130}$$

where h_{ij} is the metric on S_{D-2}, $t_i = \delta_i^t$, $k_i = \delta_i^r$, q and c_0 are constants. Then the antisymmteric tesor F_{ij} defined in (21) becomes

$$F_{ij} = \frac{c_0}{r^2}(t_i k_j - t_j k_i) \tag{131}$$

$$\widetilde{M_{ij}} = F_{mj} F_i^m - \frac{1}{4} F^2\, g_{ij} \tag{132}$$

$e_1 = e_3 = 0 \Rightarrow\ S_{ij} = e_0 \widetilde{M_{ij}},\ e_0 = q^2$

$$R_{ijkl} = q^2 [g_{jl} \widetilde{M_{ik}} - g_{jk} \widetilde{M_{il}} + g_{ik} \widetilde{M_{jl}} - g_{il} \widetilde{M_{kj}}] + \frac{q^2(c_0^2 - 1)}{c_0^2} F_{ij} F_{ml} \tag{133}$$

$$R_{ij} = q^2[(D-3) + \frac{1}{c_0^2}] \widetilde{M_{ij}} + \frac{1}{2q^2}[(D-3) - \frac{1}{c_0^2}] g_{ij} \tag{134}$$

$$G_{ij} = q^2[(D-3) + \frac{1}{c_0^2}] \widetilde{M_{ij}} + \frac{1}{2q^2}[\frac{1}{c_0^2} - (D-3)^2] g_{ij} \tag{135}$$

It is easily seen that

$$\nabla_l F_{ij} = 0\ ,\ \nabla_m R_{ijkl} = 0 \tag{136}$$

Here the origin of the antisymmteric tensor F_{ij} is geometrical. It is the volume form of the two dimensional part $ds^2 = -A^2\, dt^2 + B^2\, dr^2$ of the D-dimesional spacetime line element (1). We can define the electromagnetic tensor field , F_{ij}^{el} , simply as $F_{ij}^{el} = Q\, F_{ij}$. Here Q is a constant (in asymptotically flat cases Q plays the role of electric charge). Eq.(135) becomes

$$G_{ij} = \frac{q^2}{Q^2}\, [(D-3) + \frac{1}{c_0^2}]\, M_{ij}^{el} + \frac{1}{2q^2} [\frac{1}{c_0^2} - (D-3)^2] g_{ij} \tag{137}$$

where M_{ij}^{el} $(= Q^2\ \widetilde{M_{ij}})$ is the energy momentum tensor of the electromagnetic field tensor F_{ij}^{el}.

In D dimensions LCBR metric is the solution of the Einstein Maxwell equations with a cosmological constant. This constant vanishes when $(D - 3) = \frac{1}{c_0^2}$. In this case spacetime is conformally flat only when $D = 4$. For this metric, the field equations of a most general lagrangian (for instance, the lagrangian of the low energy limit of the string theory at all orders in string tension parameter) yields

$$\begin{aligned} E_{ij} &= \sigma_2 g_{ij} + \sigma_1 M_{ij} = G_{ij}, & (138)\\ E_i &= \sigma k_i, & (139)\\ E &= \sigma', & (140) \end{aligned}$$

where σ_1, σ_2, σ and σ' are constants. This means that, Einstein's equations reduce to algebraic equations. The lagrangian of the most general theory is a scalar containing the curvature tensor , metric tensor , matter fields and their derivatives , contractions and multiple products of all orders. According to the theorem 2 given above all second rank symmetric tensors constructed out of these are expressible in terms of g_{ij} , M_{ij} , and $k_i k_j$. So whatever the theory is we have three equations (under our assumptions all the terms containing k_i drop out) for five constants. Two constants (c_0, q) come from the definition of the D-dimensional LCBR metric (130). One constant Q comes from the definition of the electromagnetic field tensor F^{el}_{ij}. The other two constants are the constant dilaton field ϕ_0 and the cosmological constant λ. In some cases λ is set to zero. In these cases as well, the number of equations are less the number of unknown constants. In the general case , under our assumptions the field equations reduce to the following algebraic equations for q, c_0, Q, ϕ_0 , and λ

$$\frac{q^2}{Q^2}\,(D-3+\frac{1}{c_0^2})=\sigma_2(q,c_0,Q,\phi_0,\lambda) \tag{141}$$

$$\frac{1}{2\,q^2}\,(-(D-3)^2+\frac{1}{c_0^2})=\sigma_4(q,c_0,Q,\phi_0,\lambda) \tag{142}$$

$$0 \;=\; \sigma(q,c_0,Q,\phi_0,\lambda) \tag{143}$$

$$0 \;=\; \sigma^{'}(q,c_0,Q,\phi_0,\lambda) \tag{144}$$

where σ_1, σ_2, σ and σ' are defined in (138)-(140). Choosing ϕ $(= \phi_0)$ and the metric function C ($= c_0$ as constants the function σ given above equations vanishes identically. Hence we are left with three algebraic equations for q, c_0, Q, ϕ_0 , and λ. These algebraic equations may or may not have solutions. For intance when we consider the Lovelock theory [26] without a cosmological contant these equations have solutions when $D = 4$ and $D > 6$. With a cosmological constant these algebraic equations have solutions for $D \geq 4$. In the case of the string theory LCBR metric is a solution of the low energy limit of the string theory. LCBR spacetime in four dimensions is also known to be an exact solution of the string theory [8] , [21]. Here we have the generalization of this result for an arbitrary value of $D > 2$. If there exists a solution of the above mentioned algebraic equations then

Theorem 5 *LCBR metric is an exact solution of the string theory at all orders*

As in the case of the LCBR spacetime [8] the symmteric spaces , $\nabla_i \, R_{jklm} = 0$ and reccurrent spaces , $\nabla_i \, R_{jklm} = l_i \, R_{jklm}$ where l_i is a (gradient) vector , play effective role in constructing exact solutions of the string theory. These spaces are in general product spaces [38]. In the case of static spherically symmetric spacetimes the only possibility to have such spaces is to choose $C = c_0$. Then D dimensional spacetime becomes $M_2 \times S^{D-2}$, where M_2 is a two dimesional geometry with metric $ds^2 = -A^2 \, dt^2 + B^2 \, dr^2$.
If we take $C = c_0$ in the metric (1), then

$$\begin{aligned} \eta_0 &= \frac{c_0^{2(D-2)}}{(D-3)!} \, \frac{1}{AB} \left(\frac{A'}{B} \right)' \\ \eta_1 &= 0 \\ \eta_2 &= \xi - (D-3)\eta_0 \\ \eta_3 &= 0 \end{aligned} \qquad (145)$$

where $\xi = \frac{c_0^{2(D-3)}}{(D-4)!}$.

$$\begin{aligned} R_{ij} &= \frac{(D-3)!}{2c_0^{2(D-2)}}\left[-\eta_0+\xi\right]g_{ij}+\left[\eta_0+\xi\right]M_{ij} \\ R &= \frac{(D-3)!}{c_0^{2(D-2)}}\left[-2\eta_0+(D-2)\xi\right] \\ G_{ij} &= -\frac{(D-3)!}{2c_0^{2(D-2)}}\left[-\eta_0+(D-3)\xi\right]g_{ij}+\left[\eta_0+\xi\right]M_{ij} \end{aligned} \tag{146}$$

$$\begin{aligned} \nabla_l H_{ij\ldots m} &= 0 \\ \nabla_i k_j &= \frac{A'}{2AB^2}\,g_{ij}+\frac{c_0^{2(D-2)}}{(D-3)!}\frac{A'}{AB^2}M_{ij}-\frac{(AB)'}{AB}k_ik_j \end{aligned} \tag{147}$$

If we let η_0 to be constant, then we have to solve the equation $\frac{1}{AB}\left(\frac{A'}{B}\right)'=\alpha$. The solution is

$$B^2=\frac{A'^2}{\alpha A^2+\beta} \tag{148}$$

where α and β are constants. Hence the metric (1) can be written as

$$ds^2=-A^2dt^2+\frac{dA^2}{\alpha A^2+\beta}+c_0^2\,d\Omega_{D-2}^2 \tag{149}$$

The case $\beta=0$ is identical with the LCBR metric. As in the case of the LCBR metric the metric constants α , c_0 and the magnetic charge satisfy three coupled (due to the theorems given before) algebraic equations. If these algebraic equations have solutions then the metric given above with nonvanishing α and β is a candidate as an exact solution of the string theory.

Above we have given the Riemann tensor in terms of the volume form of S_{D-2} (D-2)-brane field. We may also write the Riemann tensor in the form

$$R_{ijml} = g_{jl}S_{im} - g_{jm}S_{il} + g_{im}S_{jl} - g_{il}S_{mj} + e_2 F_{ij}F_{ml} \tag{150}$$

In this case

$$\begin{aligned} e_0 &= c_0^2 \\ e_1 &= 0 \\ e_2 &= -c_0^2 + c_0^2\zeta \\ e_3 &= 0 \end{aligned} \tag{151}$$

where $\zeta = \frac{c_0^2}{AB}\left(\frac{A'}{B}\right)'$

$$\begin{aligned} R_{ij} &= \frac{D-3-\zeta}{2c_0^2}\,g_{ij} + (D-3+\zeta)c_0^2\widetilde{M_{ij}} \\ R &= \frac{D^2-5D+6-2\zeta}{c_0^2} \\ G_{ij} &= \frac{\zeta-(D-3)^2}{2c_0^2}\,g_{ij} + (D-3+\zeta)c_0^2\widetilde{M_{ij}} \end{aligned} \tag{152}$$

$$\tag{153}$$

$$\begin{aligned} \nabla_l F_{ij} &= 0 \\ \nabla_i k_j &= \frac{A'}{2AB^2}\,g_{ij} + c_0^4\frac{A'}{AB^2}\widetilde{M_{ij}} - \frac{(AB)'}{AB}k_ik_j \end{aligned} \tag{154}$$

The solution given in (148) is not valid when A is set to a constant. In such a case B can be choosen as unity. Therefore letting $A = B = 1$ and $C = c_0$. We find that $e_0 = -e_2 = c_0^2$ and

$$\nabla_l F_{ij} = 0 \ , \ \nabla_m R_{ijkl} = 0 \tag{155}$$

$$\nabla_i k_j = 0 \ , \ \nabla_i t_j = 0 \tag{156}$$

$$R_{ij} = (D-3)c_0^2 \left[\widetilde{M}_{ij} + \frac{1}{2c_0^2} g_{ij}\right] \tag{157}$$

$$R = \frac{(D-3)(D-2)}{c_0^2} \tag{158}$$

$$G_{ij} = (D-3)c_0^2 \left[\widetilde{M}_{ij} + \frac{1}{2c_0^2}(3-D) g_{ij}\right] \tag{159}$$

$$\tag{160}$$

Since the metric contains only one arbitrary constant c_0 it may not be possible to eliminate the coefficient of g_{ij} (the cosmological constant) in the field equations. Hence the metric given by

$$ds^2 = -dt^2 + dr^2 + c_0^2 \, d\,\Omega_{D-2}^2 \tag{161}$$

is a canditate as an exact solution of the string equations with a cosmological constant.

6 Conclusion

We have found all possible asymptotically flat solutions of Einstein Maxwell Dilaton field theory in D dimensional static spherically symmetric spacetimes with arbitrary dilaton coupling constant. We proved that the asymptotically flat black hole solutions carrying mass M , electric charge Q and dilaton charge Σ in four dimensions are unique not only in static spherically symmteric spacetimes but also in static axially symmetric spacetimes. This proof may be extended to arbitrary dimensions and also to spacetimes which are not asymptotically flat [24]. We searched for all possible static spherically spacetimes which may solve the string equations exactly (to all orders in string tension parameter).

This work is partially supported by the Scientific and Technical Research Council of Turkey (TUBITAK) and Turkish Academy of Sciences (TUBA). We would like to thank the referees for their several suggestions , in particular for informing us the recent works [23] , [24].

References

[1] A. A. Tseytlin , *Exact string solutions and duality* , To appear in: Proceedings of the 2nd Journe'e Cosmologie, Observatoire de Paris , June 2-4 ,1994 (World Scientific, Singapore) Imperial/TP/93-94/46 , hep-th/9407099

[2] R. Güven , Phys. Lett. **B191** , 275 (1987).

[3] D. Amati and C. Klimcik , Phys. Lett. **B219** , 443 (1989).

[4] G. Horowitz and A. Steif , Phys. Rev. Lett. **64** , 260 (1990).

[5] G. Horowitz and A. A. Tseytlin , Phys. Rev. Lett. **73** , 3351 (1994).

[6] A. A. Tseytlin , *Black holes and Exact Solutions in String Theory* , to appear in the proceedings of **International School of Astrophysics "D. Chalonge" , 3rd course: Current Topics in Astrofundamental Physics** , 4-16 September 1994 , Erice , ed. N. Sanchez (World Scientific , Singapore).

[7] T. Witten and Nappi , Phys. Rev. Lett. **71** , 3751 (1993).

[8] D. Lowe and A. Strominger , Phys. Rev. Lett. **73** , 1468 (1994)

[9] R. Kallosh and T. Ortin , Phys.Rev.**D50** R7123 (1994).

[10] G. Horowitz and A. A. Tseytlin , Phys.Rev **D51** , 2896 (1995).

[11] Barton Zwiebach, Phys. Lett. **156B** (1985) 315.

[12] David G. Boulware and S. Deser, Phys. Rev. Lett. **55** (1985) 2656.

[13] Bruno Zumino, Phys. Reports **137** (1986) 109.

[14] D. L. Wiltshire, Phys. Lett. 168B (1986) 36.

[15] James T. Wheeler, Nucl. Phys. **B273** (1986) 732.

[16] Brian Whitt, Phys. Rev. **D38** (1988) 3000.

[17] G.W. Gibbons and K. Maeda , Nucl.Phys.**B298**, (1988) 741.

[18] D. Garfinkle , G.T. Horowitz and A. Strominger , Phys.Rev.**D 43** , (1991) 3140.

[19] M.J. Duff and J.X. Lu , Nuclear Physics **B416** , 301 (1994)

[20] M.J. Duff, R.R. Khuri and J.X. Lu , *String Solitons* , preprint , NI-94-017, CTP/TAMU-67/92 , McGill/94-53 , CERN-TH.7542/94. hep-th/9412184.

[21] M. Gürses , Phys.Rev.**D 46** (1992) 2522.

[22] M. Rakhmanov , Phys. Rev. **D50** 5155 (1994).

[23] S. J. Poletti and D.L. Wiltshire , Phys. Rev. **D50** 7260 (1994).

[24] Kevin C.K. Chan , James H. Horne , and Robert Mann , *Charged Dilaton Black Holes with Unusal Asymptotics* , preprint , DAMTP/P-95/7 , WATPHYS TH-95/02 (gr-qc/9502042).

[25] G.W. Gibbons and C.G. Wells , *Anti-Gravity Bounds and the Ricci Tensor* (DAMTP preprint R93/25) , (gr-qc/9310002)

[26] D. Lovelock , J. Math. Phys. **12**, 498 (1971) ; **13**, 874 (1972).

[27] A. K. M. Masood-ul-Alam , Class. Quantum Grav. **10** 2649 (1993).

[28] G. Bunting , *The Black Hole Uniqueness Theorem* , in **Proc. of the and Australian Mathematics Convention** , Sydney ,(1981).

[29] G. Bunting , *Proof of the Uniqueness Conjecture for Black Holes* , Ph.D. Thesis , University of New England , Armidale ,N.S.W. (1983).

[30] P.O. Mazur , J.Phys. **A15** , 3173 (1982).

[31] P.O. Mazur , Acta. Phys. Pol. **B14** , 219 (1983).

[32] B. Carter , Commun. Math. Phys. **99** , 563 (1985).

[33] P. Breitenlohner , D. Maison and G. Gibbons , Commun. Math. Phys. **120** 295 (1988).

[34] M. Gürses , Letters in Mathematical Physics , **16** 265 (1992).

[35] M. Gürses and B. Xanthopoulos , Phys.Rev.**26D** , 1912 (1982).

[36] M. Gürses , Phys.Rev. **30D** ,486 (1984).

[37] A. Eriş , M. Gürses and A. Karasu , J. Math. Phys. **25** , 1489 (1984).

[38] H.S.Ruse , A.G.Walker and T.J.Willmore , *Harmonic Spaces* , Consiglio Nazionale Delle Richerche Monografie Matematiche No.8, (Edizioni Cremonese Roma).